On Heterotic Noncompact Nonlinear Sigma Models

Peter Koroteev

University of Minnesota



In collaboration with A. Monin, M. Shifman, W. Vinci, A. Yung

1009.6207 1107.3779 work in progress

SUSY 2011 Fermilab IL August 28th

Outline

- N=2 SQCD in 4d & Sigma models in 2d
- GLSM as a tool to find BPS spectrum
- Heterotic deformation and Large-N solution - beyond BPS sector
- Modeling confinement
- Perturbation theory

4d SQCD vs 2d sigma models

4d / 2d duality

$$\mathcal{N} = 2 \quad SU(N) \quad \mathbf{SQCD}$$

 $N_f = N + \tilde{N}$ fund hypers w/ masses

$$m_1, \dots, m_N \quad \mu_1, \dots, \mu_{\tilde{N}}$$

$$\tau = \frac{4\pi i}{g^2} + \frac{\theta}{2\pi}$$

on baryonic Higgs branch

(2,2) U(1) GLSM e

N chiral + I \tilde{N} chiral - I w/ twisted masses

$$m_1, \dots, m_N \qquad \mu_1, \dots, \mu_{\tilde{N}}$$

$$\tau = ir + \frac{\theta}{2\pi}$$

vortex moduli space


BPS dyons (Seiberg-Witten)


kinks interpolating between different vacua

BPS spectra (as functions of masses, Lambda) are the same

Brane construction

[Hanany Tong]





D-term

take k=Ito get U(I) theory on DI (semilocal vortex)

$$|Q_i|^2 - |\tilde{Q}_j|^2 = r$$

FI term -- separation of NS5s in x6

will refer to as HT model

$U(N_c)$ $\mathcal{N}=2$ d=4 SQCD w/ N_f quarks

$$\{Q_{\alpha}^{I},\bar{Q}_{\dot{\beta}}^{J}\}=2\delta^{IJ}P_{\alpha\dot{\beta}}+2\delta^{IJ}Z_{\alpha\dot{\beta}}$$

$$\{Q_{\alpha}^{I},Q_{\beta}^{J}\}=2Z_{\alpha\beta}^{IJ}$$
 monopoles

$$\mathcal{L} = \operatorname{Im} \left[\tau \int d^{4}\theta \operatorname{Tr} \left(Q^{i\dagger} e^{V} Q_{i} + \tilde{Q}^{i\dagger} e^{V} \tilde{Q}_{i} + \Phi^{\dagger} e^{V} \Phi \right) \right] + \operatorname{Im} \left[\tau \int d^{2}\theta \left(\operatorname{Tr} W^{\alpha 2} + m_{j}^{i} \tilde{Q}_{i} Q^{j} + Q_{i} \Phi \tilde{Q}^{i} \right) \right]$$

bosonic part

$$S = \int d^4x \operatorname{Tr} \left\{ \frac{1}{2g^2} F_{\mu\nu}^2 + \frac{1}{g^2} |D_{\mu}\Phi|^2 + |\nabla_{\mu}Q|^2 + \frac{g^2}{4} (Q\bar{Q} - \xi)^2 + |\Phi Q + QM|^2 \right\}$$

BPS conditions

$$B_3 - g^2(Q\bar{Q} - \xi^2) = 0$$
$$\nabla_3 Q = 0$$

String tension

$$T = \xi \int d^2x \operatorname{Tr} F_{12} = 2\pi \xi n$$

Vortex moduli space

Nf=Nc color-flavor locked phase single SUSY vacuum

$$U(N_c) \times SU(N_f) \to SU(N)$$


local vortex

$$\frac{SU(N)}{SU(N-1)\times U(1)} = \mathbb{CP}^{N-1}$$


Nf>Nc semilocal

$$\pi_2(\mathcal{M}_{vac}) = \pi_2 \left(\frac{SU(N+\tilde{N})}{SU(N) \times SU(\tilde{N}) \times U(1)} \right) = \mathbb{Z}$$

Duality between two strongly coupled theories



Confined monopoles



Hanany-Tong model as U(I) GLSM

$$\mathcal{L} = \int d^4\theta \left[\sum_{i=1}^{N_c} \Phi_i^{\dagger} e^{\mathcal{V}} \Phi_i + \sum_{i=1}^{\tilde{N}} \widetilde{\Phi}_i^{\dagger} e^{-\mathcal{V}} \widetilde{\Phi}_i - r\mathcal{V} + \frac{1}{2e^2} \Sigma^{\dagger} \Sigma \right]$$

Limit $e o \infty$ defines vacuum manifold $\mathcal{O}(-1)^{\otimes \tilde{N}}$ over \mathbb{CP}^{N-1}

 $R_{i\bar{\jmath}} = \frac{N-N}{m} g_{i\bar{\jmath}} + \mathcal{O}(\frac{1}{m^2})$

perturbation theory is subtle

$$\beta_{i\bar{\jmath}} = a^{(1)} R_{i\bar{\jmath}} \log \frac{M}{\mu} + \dots$$

however there are nonperturbative corrections

One loop twisted effective superpotential is exact in (2,2)

$$\widetilde{\mathcal{W}}_{eff} = \frac{N - \widetilde{N}}{4\pi} \sigma \left(\log \frac{\sigma}{\Lambda} - 1 \right)$$

$$V = \theta^{+}\bar{\theta}^{+}(A_{0} + A_{3}) + \theta^{-}\bar{\theta}^{-}(A_{0} - A_{3}) - \theta^{-}\bar{\theta}^{+}\sigma - \theta^{-}\bar{\theta}^{+}\bar{\sigma} + \bar{\theta}^{2}\theta\lambda + \theta^{2}\bar{\theta}\bar{\lambda} + \bar{\theta}\theta\bar{\theta}\theta D$$

gives vacua of the theory and its BPS spectrum !!

Heterotic deformation

(0,2) Theory

In 4d introduce masses

breaks
$$\mathcal{N}=2$$
 to $\mathcal{N}=1$

obtain heterotic sigma model

$$\int d^2\theta \, \mu^2 (\Phi^a)^2$$

On the flux tube $(2,2) \longmapsto (0,2)$

[Edalati Tong][Shifman Yung] [Distler Kachru]

$$\mathcal{L} = \int d^4\theta \left(\Phi_i^{\dagger} e^V \Phi^i - rV - \mathcal{B}V \right) \qquad \mathbb{CP}^{N-1} \times \mathbb{C}$$

B-right handed superfield

can be treated as model w/ field dependent FI term

$$K = (r + \mathcal{B})\log(1 + |\phi^i|^2)$$

Geometry becomes non-Kahler

due to generation of H field (field dependent theta term)

(0,2) GLSM

$$\int d^4\theta \left[\sum_{i=1}^{N_c} \Phi_i^{\dagger} e^V \Phi_i + \sum_{i=1}^{N_c - N_f} \tilde{\Phi}_i^{\dagger} e^{-V} \tilde{\Phi}_i - (r + \mathcal{B})V + \frac{1}{2e^2} \Sigma^{\dagger} \Sigma \right]$$

$$\Phi^{i} = n^{i} + \bar{\theta}\xi^{i} + \theta\bar{\xi}^{i} + \bar{\theta}\theta F^{i}, \quad i = 1, \dots, N_{c}$$

$$\widetilde{\Phi}^{j} = \rho^{j} + \bar{\theta}\eta^{j} + \theta\bar{\eta}^{j} + \bar{\theta}\theta\tilde{F}^{j}, \quad j = 1, \dots, \tilde{N}$$

$$\Sigma = \sigma + i\theta^{+}\bar{\lambda}_{+} - i\bar{\theta}^{-}\lambda_{-} + \theta^{+}\bar{\theta}^{-}(D - iF_{01})$$

$$\mathcal{B} = \omega(\bar{\theta}\zeta_R + \bar{\theta}\theta\bar{\mathcal{F}}\mathcal{F})$$

deformation adds

$$\mathcal{L}^{het} = \mathcal{L} + \bar{\zeta}_R \partial_L \zeta_R - |\omega|^2 |\sigma|^2 - [i\omega \lambda_L \zeta_R + \text{H.c.}]$$


Not enough SUSY to fix superpotential Have to dwell on large-N approach

Large-N solution of (0,2)

$$V_{1-loop} = \frac{1}{4\pi} \sum_{i=1}^{N-1} \left(-\left(D + |\sigma - m_i|^2\right) \log \frac{|\sigma - m_i|^2 + D}{\Lambda^2} + |\sigma - m_i|^2 \log \frac{|\sigma - m_i|^2}{\Lambda^2} \right)$$
$$- \frac{1}{4\pi} \sum_{j=1}^{\tilde{N}-1} \left(-\left(D - |\sigma - \mu_j|^2\right) \log \frac{|\sigma - \mu_j|^2 - D}{\Lambda^2} - |\sigma - \mu_j|^2 \log \frac{|\sigma - \mu_j|^2}{\Lambda^2} \right)$$
$$+ \frac{N - \tilde{N}}{4\pi} D.$$

$$V_{eff} = V_{1-loop} + (|\sigma - m_0|^2 + D) |n_0|^2 + (|\sigma - \mu_0|^2 - D) |\rho_0|^2 + \frac{uN}{4\pi} |\sigma|^2$$

for zero masses



Symmetric masses

$$m_k = m e^{2\pi i \frac{k}{N}}, \quad k = 0, \dots, N - 1,$$

 $\mu_l = \mu e^{2\pi i \frac{l}{N}}, \quad l = 0, \dots, \tilde{N} - 1.$

Vacuum equations

$$(|\sigma - m_0|^2 + D) n_0 = 0, \quad (|\sigma - \mu_0|^2 - D) \rho_0 = 0,$$

$$\frac{1}{4\pi} \sum_{i=1}^{N-1} \log \frac{|\sigma - m_i|^2 + D}{\Lambda^2} - \frac{1}{4\pi} \sum_{j=1}^{\tilde{N}-1} \log \frac{|\sigma - \mu_j|^2 - D}{\Lambda^2} = |n_0|^2 - |\rho_0|^2,$$

$$\frac{1}{4\pi} \sum_{i=1}^{N-1} (\sigma - m_i) \log \frac{|\sigma - m_i|^2 + D}{|\sigma - m_i|^2} + \frac{1}{4\pi} \sum_{j=1}^{\tilde{N}-1} (\sigma - \mu_j) \log \frac{|\sigma - \mu_j|^2 - D}{|\sigma - \mu_j|^2} =$$

$$= (\sigma - m_0) |n_0|^2 + (\sigma - \mu_0) |\rho_0|^2 + \frac{uN}{4\pi} \sigma.$$

Solution of (2,2) model

Phase transitions -- artifact of large-N

$$(|\sigma - m_0|^2 + D) n_0 = 0, \quad (|\sigma - \mu_0|^2 - D) \rho_0 = 0$$

Higgs in n (Hn)

$$\rho_0 = 0 \quad D = -|\sigma - m|^2$$

Higgs in rho (Hrho)

$$n_0 = 0 \quad D = |\sigma - \mu|^2$$

Coulomb (C)

$$r = \left\{ \begin{array}{ll} \frac{N-\tilde{N}}{2\pi}\log\frac{m}{\Lambda}\,, & \mu < m \\ \\ \frac{N}{2\pi}\log\frac{m}{\Lambda} - \frac{\tilde{N}}{2\pi}\log\frac{\mu}{\Lambda}\,, & \mu > m \,. \end{array} \right.$$

$$r = \begin{cases} \frac{N - \tilde{N}}{2\pi} \log \frac{\mu}{\Lambda}, & \mu > m \\ \\ \frac{N}{2\pi} \log \frac{m}{\Lambda} - \frac{\tilde{N}}{2\pi} \log \frac{\mu}{\Lambda}, & \mu < m \end{cases}$$


$$n_0 = \rho_0 = 0$$

renormalized FI term vanishes in C phase


in (2,2) from exact superpotential

$$\frac{\prod_{i}(\sigma - m_i)}{\prod_{i}(\sigma - \mu_j)} = \Lambda^{N - \tilde{N}} \qquad \sigma = 0$$


is one of the solutions...




N=5 Nf=8




N=15 Nf=18



Nf=5 C μ phase







Spectrum

[Bolokhov Shifman Yung]
[PK Monin Vinci]

$$\mathcal{L} = -\frac{1}{4e_{\gamma}^2} F_{\mu\nu}^2 + \frac{1}{e_{\sigma 1}^2} (\partial_{\mu} \Re \sigma)^2 + \frac{1}{e_{\sigma 2}^2} (\partial_{\mu} \Im \sigma)^2 + i \Im (\bar{b} \, \delta \sigma) \epsilon_{\mu\nu} F^{\mu\nu} - V_{\text{eff}}(\sigma) + \text{Fermions}$$

Anomaly





$$b = \frac{N}{4\pi} \left(\frac{1}{N} \sum_{i=1}^{N-1} \frac{1}{\bar{\sigma}_0 - \bar{m}_i} - \alpha \frac{1}{\tilde{N}} \sum_{i=1}^{\tilde{N}-1} \frac{1}{\bar{\sigma}_0 - \bar{\mu}_i} \right)$$

$$m_{\gamma} = e_{\sigma \, 2} e_{\gamma} |b|$$

Photon becomes massless in Cs phase!! Confin

Confinement!

Note that Lambda vacua disappear at large deformations

Need to sit in zero-vacua

e.g. in Cm phase

$$m_{\gamma} = \sqrt{6} \Lambda \left(\frac{\Lambda}{m}\right)^{1/\alpha} \left(\left(\frac{m}{\Lambda}\right)^{2/\alpha} - \left(\frac{\mu}{\Lambda}\right)^2 e^{u/\alpha}\right) e^{-\frac{u}{2\alpha}}$$

Massless goldstino in fermionic sector

Conclusions and open questions

- Study BPS (and beyond) spectrum of SQCD can effectively be done using 2d NLSM (and GLSM)
- Rich variety of phases in (0,2) model at strong coupling
- ullet Other heterotic deformations $ar{D}\Phi_+\simar{D}\Phi_-$
- Are there flux tubes in theories without FI term?
 (e.g. SU(N)) Omega deformed 4d theory may have such solutions...
- Connections to integrable systems in 2d...
- Relationship w/ another 4d/2d duality [Vafa et al]